

Appendix I: Using Summation Notation With GAMS	AI-1
AI.1 Summation Mechanics.....	AI-1
AI.1.1 Sum of an Item.....	AI-1
AI.1.2 Multiple Sums.....	AI-2
AI.1.3 Sum of Two Items	AI-2
AI.2 Summation Notation Rules.....	AI-3
AI.2.1 For a Scalar Equation	AI-3
AI.2.2. For a Family of Equations	AI-4
AI.4 Defining and Using Variables	AI-7
AI.5 Equations	AI-8
AI.6 Cautions and Extensions.....	AI-9

Appendix I: Using Summation Notation With GAMS

Summation notation is difficult for some students to use and follow. Here we present notes on the mechanics of summation notation usage and some rules for proper use. This discussion is cast within the GAMS framework with presentation equivalents of common summation expressions and error messages caused by improper summation. All of the GAMS statements used herein are shown in Table 1 and are in file NOTATION.

AI.1 Summation Mechanics

Summation notation is a short hand way of expressing sums of algebraic terms involving subscripted items. In order to cover the mechanics of summation notation it is useful to have a set of subscripted items and associated numerical values. Thus, let us define some data

$$\begin{array}{l} x_1 = 1 \quad y_{11} = 2 \quad y_{12} = 3 \\ x_2 = 2 \quad y_{21} = 4 \quad y_{22} = 1 \\ x_3 = 3 \quad y_{31} = 1 \quad y_{32} = 4. \end{array}$$

Now let us define a variety of summation expressions.

AI.1.1 Sum of an Item.

Suppose we wished to sum all values of x . This would be written as

$$\sum_{i=1}^3 x_i = x_1 + x_2 + x_3 = 1 + 2 + 3 = 6$$

or in GAMS

$$\text{SUM1} = \text{SUM}(I, X(I));$$

For short hand purposes if i was to be summed over all possible values, we would write this as

$$\sum_i x_i .$$

We might also express a sum as follows which indicates all of the i are summed over except $i=3$

$$\sum_{i=1}^3 x_i = 3.$$

In GAMS, this is more difficult to express where one has to write a conditional (\$) operation or define a subset as follows

$$\text{SUM1} = \text{SUM}(\text{I}\$(\text{ORD}(\text{I.NE.3})), \text{X}(\text{I}));$$

or

$$\begin{aligned} \text{SET SUBSETI}(\text{I}) &= /1, 2/; \\ \text{SUM1} &= \text{SUM}(\text{SUBSETI}, \text{X}(\text{SUBSETI}(\text{I}))); \end{aligned}$$

AI.1.2 Multiple Sums

Sums over two indices consider all combinations of those items

$$\sum_i \sum_j y_{ij} = y_{11} + y_{12} + y_{21} + y_{22} + y_{31} + y_{32} = 15.$$

The equivalent GAMS expression is

$$\text{SUM2} = \text{SUM}(\text{I,J}, \text{Y}(\text{I,J}));$$

AI.1.3 Sum of Two Items

Suppose we wished to sum over two items completely where they shared a subscript

$$\sum_{i=1}^3 (x_i + \sum_{j=1}^2 y_{ij}) = \sum_i (x_i + \sum_j y_{ij}) = \sum_i x_i + \sum_i \sum_j y_{ij} = x_1 + y_{11} + y_{12} + x_2 + y_{21} + y_{22} + x_3 + y_{31} + y_{32} = 21.$$

The equivalent GAMS expression is as follows

$$\text{SUM3} = \text{SUM}(\text{I}, \text{X}(\text{I}) + \text{SUM}(\text{J}, \text{Y}(\text{I}, \text{J})));$$

or

$$\text{SUM3} = \text{SUM}(\text{I}, \text{X}(\text{I})) + \text{SUM}(\text{I,J}, \text{Y}(\text{I,J}));$$

On the other hand, if we wished to sum the results only for the i^{th} element and call it A_i then

$$A_i = x_i + \sum_j y_{ij} = x_i + y_{i,1} + y_{i,2} \text{ or in GAMS}$$

$$A(\text{I}) = \text{X}(\text{I}) + \text{SUM}(\text{J}, \text{Y}(\text{I,J}));$$

which would yield a vector [6 , 7 , 8] of results.

Sums over common subscripts can be collapsed or taken apart

$$\sum_i (x_i + z_i) = \sum_i x_i + \sum_i z_i \quad \text{or}$$

$$\text{SUM4} = \text{SUM}(I, X(I) + Z(I));$$

or

$$\text{SUM4} = \text{SUM}(I, X(I)) + \text{SUM}(I, Z(I));$$

AI.2 Summation Notation Rules

Certain rules apply when writing summation notation equations. The applicable rules depend on whether the final result is an unsubscripted scalar or a subscripted family of results determined by multiple equations.

AI.2.1 For a Scalar Equation

$$B1 = \sum_i \sum_j \sum_k p_{ijk} + \sum_m \sum_n q_{submn}.$$

All subscripts must be dealt with in each term. Thus, it is proper to define the equation

However, the following equations are wrong

$$B2 = p_{ijk} + q_{mn}$$

$$B3 = \sum_j \sum_i p_{ijk} + \sum_m \sum_n q_{mn}.$$

In the case of the first equation, the result would really have the subscripts i,j,k,m,n,

while the second equation result would have to have a k subscript on B3 or a sum over k to be proper.

Equivalent GAMS commands for the above equation expressions are

$$\text{EQB1.. B1 =E= SUM}((I,J,K),P(I,J,K)) + \text{SUM}((M,N), Q(M,N));$$

$$\text{EQB2.. B2 =E= P(I,J,K) + Q(M,N);}$$

$$\text{EQB3.. B3 =E= SUM}((I,J), P(I,J,K)) + \text{SUM}((M,N), Q(M,N));$$

Here, the first equation expression is correct, while the last two equation expressions are incorrect. If you run GAMS with the above commands, you would encounter GAMS error messages \$149 which says

"UNCONTROLLED SET ENTERED AS CONSTANT" meaning that you have not somehow dealt with all the subscripts in the equation.

AI.2.2. For a Family of Equations

Several rules apply when one is working with a family of equations.

1. The members of the family must be specified with an indication of the subscripts which define each equation. This is done by indicating all the conditions for which the equations exist in a "for" condition. For example, suppose we define an equation which sets all C's equal to 2. This is done by saying

$$C_i = 2 \quad \text{for all } i \quad \text{or} \quad C_i = 2 \quad \text{for } i = 1, 2, \dots, n.$$

Similarly, if we wish to set a 2 dimensional variable equal to a constant, we would state

$$D_{ij} = 2 \quad \text{for all } i \text{ and } j,$$

while stating that for each row of the matrix E_{ij} we have the same values F_i is defined by

$$E_{1ij} = F_i \quad \text{for all } i \text{ and } j.$$

The equivalent GAMS commands for the above expressions are

```

EQUATIONS
      EQC(I)          EQUATION C
      EQD(I,J)        EQUATION D
      EQE1(I,J)       EQUATION E1;
EQC(I).. C(I) =E= 2;
EQD(I,J).. D(I,J) =E= 2;
EQE1(I,J).. E1(I,J) =E= F(I);

```

On the other hand, it is wrong to state

$$E_{2ij} = 2$$

without conditions on i and j . The equivalent GAMS commands for the above incorrect expressions are

```

EQUATION
      EQE2          EQUATION E2;
EQE2.. E2(I,J) =E= 2;

```

Here you would get error message \$149 which says "UNCONTROLLED SET

ENTERED AS CONSTANT."

2. When writing an equation with a for statement, all subscripts which are not in the for

$$\sum_j \sum_k p_{ijk} = G1_i \quad \text{for all } i$$

$$\sum_k p_{ijk} = H1_i \quad \text{for all } i \text{ and } j$$

statement must be summed over. Consequently, it is proper to write

but improper to write

$$p_{ijk} = G2_i \quad \text{for all } i$$

$$\sum_k p_{ijk} = H2_i \quad \text{for all } i.$$

The equivalent GAMS commands for the above equations are

```
EQUATIONS
    EQG1(I) EQUATION G1
    EQH1(I,J) EQUATION H1
    EQG2(I) EQUATION G2
    EQH2(I) EQUATION H2;
EQG1(I).. G1(I) =E= SUM((J,K), P(I,J,K));
EQH1(I,J).. H1(I,J) =E= SUM(K, P(I,J,K));
EQG2(I).. G2(I) =E= P(I,J,K);
EQH2(I).. H2(I) =E= SUM(K, P(I,J,K));
```

in which the first two equations are correct, while the last two equations are wrong and error messages \$149 "UNCONTROLLED SET ENTERED AS CONSTANT" would again be realized.

3. In any term of an equation, the result after executing the mathematical operations in that term must be of a dimension less than or equal to the family definition in the for statement. For example, it is proper to write

$$\sum_j \sum_k p_{ijk} = L1 \quad \text{for all } i$$

$$\sum_j \sum_k r_{ijkm} + \sum_j s_{ijm} = N_{im} \quad \text{for all } i \text{ and } m$$

but wrong to write

$$p_{ijk} = L2 \quad \text{for all } i.$$

Thus, for the following expressions, the first two equations are appropriate but the last equation would give you error message \$149 "UNCONTROLLED SET ENTERED AS CONSTANT."

```

EQUATION
EQLI(I)...    LI(I) =E=    SUM((J,K), P(IJK));
EQNI(I,M)... N(I,M) =E=    SUM((J,K), R(I,J,K,M))
              + SUM (J,S(I,J,M));
EQL2(I)...    L2 =E=    P(I,J,K);

```

4. When the dimension is less than the family definition this implies the same term appears in multiple equations. For example, in the equation

$$2 + \sum_j \sum_k p_{ijk} + \sum_j s_{ijm} = O_{im} \quad \text{for all } i \text{ and } m,$$

the 2 term appears in every equation and the sum involving p is common when m varies.

Equivalent GAMS commands are as follows

```

EQUATION
EQO(I,M)    EQUATION O;
EQO(I,M)..  2 + SUM((J,K), P(I,J,K)) + SUM(J, S(I,J,M)) =E= O(I,M);

```

5. In an equation you can never sum over the parameter that determines the family of equations. It is certainly wrong to write

$$\sum_k \sum_j \sum_i p_{ijk} = W_i \quad \text{for all } i.$$

Or, equivalently, the following expressions are wrong and will result in error

message \$125 which says "SET IS UNDER CONTROL ALREADY."

```
EQW(I)...      W(I) =E=      SUM(I,J,K), P(I,J,K);
```

AI.3 Defining Subscripts

In setting up a set of equations and variables use the following principles. Define a subscript for each physical phenomena set which has multiple members, i.e.,

Let i denote production processes of which there are I
 j denote locations of which there are J
 k denote products of which there are K
 m denote sales locations of which there are M.

Equivalent GAMS commands are

```
SET    I        /1*20/  
      J        /1*30/  
      K        /1*5/  
      M        /CHICAGO, BOSTON/;
```

Define different subscripts when you are either considering subsets of the subscript set or different physical phenomena.

AI.4 Defining and Using Variables

1. Define a unique symbol with a subscript for each manipulatable item.

For example:

p_{ijk} = production using process i at location j while producing good k.

Or, equivalently,

```
PARAMETER   P(I,J,K)
```

or

```
PARAMETER   PRODUCTION(PROCESS, LOCATION, GOOD)
```

Here, for documentation purposes, the second expression is preferred.

2. Make sure that variable has the same subscript in each place it occurs.

Thus it is proper to write

$$\begin{aligned} \text{Max } & \sum_i \sum_j \sum_k t_{ijk} \\ & \sum_i \sum_j t_{ijk} = 3 \quad \text{for all } k \end{aligned}$$

but wrong to write

$$\begin{aligned} \text{Max } & \sum_i \sum_j t_{ij} \\ & \sum_i \sum_j t_{ijk} = 3 \quad \text{for all } k \\ & t_{ijk} \geq 0. \end{aligned}$$

The second model would cause error message \$148 indicating "DIMENSION DIFFERENT."

3. The authors feel it is a bad practice to define different items with the same symbol but varying subscripts. We think you should never use the same symbol for two different items as follows

u_{ij} = amount of tires transported from i to j and

u_{kj} = amount of chickens transported from k to j .

GAMS would not permit this, giving error \$150 "Symbolic Equations Redefined."

AI.5 Equations

Modelers should carefully identify the conditions under which each equation exists and use subscripts to identify those conditions. We do not think modelers should try to overly compact the families of equations. For example, it is OK to define

$$\sum_j a_{ij} x_j \# b_i$$

for all i , where a_{ij} is use of water by period and labor by period, where i denotes water periods and labor periods and b_i simultaneously contains water and labor availability by period. But we find it is better to define

$$\sum_j d_{ij} x_j \# e_i$$

$$\sum_j f_{ij} x_j \# h_i$$

where i denotes period,

d_{ij} denotes water use and e_i water availability,

f_{ij} denotes labor use and h_i labor availability.

AI.6 Cautions and Extensions

1. Be careful when you sum over terms which do not contain the subscript you are summing over. This is equivalent to multiplying a term by the number of items in the sum.

$$\sum_{j=1}^N x_i = Nx_i$$

$$\sum_{j=1}^3 X_2 = 3(2) = 6$$

Or, in GAMS

$$\text{SUM5A} = \text{SUM}(J, X("2"));$$

2. Be careful when you have a term in a family of equations which is of a lesser dimension than the family, this term will occur in each equation. For example, the expression

$$\sum_j x_j = z_i \quad \text{for } i = 1,2,3$$

implies that simultaneously

$$\sum_j x_j = z_1$$

$$\sum_j x_j = z_2$$

$$\sum_j x_j = z_3.$$

3. The same rules as outlined above apply to product cases

$$\prod_{i=1}^3 x_i = x_1 * x_2 * x_3.$$

Or, equivalently,

$$\text{PRODUCTX} = \text{PROD}(\text{I}, \text{X}(\text{I}));$$

4. The following relationships also hold for summation

a. $\sum_i K x_i = K \sum_i x_i$

b. $\sum_{i=1}^n KP = K \sum_{i=1}^n P = K n P$

c. $\sum_i \sum_j (v_{ij} + y_{ij}) = \sum_i \sum_j v_{ij} + \sum_i \sum_j y_{ij}$

d. $\sum_i \sum_j (x_i + y_{ij}) = n \sum_i x_i + \sum_i \sum_j y_{ij}$ when $j = 1, 2, \dots, n$

Table 1. Sample GAMS Commands for Summation Notation Expressions

```

1 *****
2 ** THIS FILE CONTAINS GAMS EXAMPLES IN SUPPORT **
3 ** OF THE NOTES USING THE SUMMATION NOTATION **
4 *****
5
6 SETS
7     I /1*3/
8     J /1*2/
9     K /1*2/
10    M /1*2/
11    N /1*3/
12
13 PARAMETERS
14
15     X(I) /1 1,2 2,3 3/
16     Z(I) /1 2,2 4,3 6/
17
18 TABLE Y(I,J)
19
20         1  2
21     1    2  3
22     2    4  1
23     3    1  4;
24
25 TABLE V(I,J)
26
27         1  2
28     1    2  3
29     2    4  1
30     3    1  4;
31
32 TABLE P(I, J, K)
33
34         1.1  1.2  2.1  2.2
35     1    1   3   5   7
36     2    2   4   6   8
37     3    1   2   3   4 ;
38
39 TABLE Q(M, N)
40
41         1  2  3
42     1    1  5 10
43     2   10  5  1;
44
45 *****
46 ** AI.1.1 SUM OF AN ITEM **
47 *****
48
49 PARAMETER
50     SUM1      SUM OF AN ITEM;
51     SUM1      = SUM(I, X(I));
52     DISPLAY SUM1;
53
54 *****
55 ** AI.1.2 MULTIPLE SUMS **
56 *****
57
58 PARAMETER
59     SUM2      MULTIPLE SUMS;
60     SUM2      = SUM((I,J), Y(I,J));
61     DISPLAY SUM2;

```

Table 1. Sample GAMS Commands for Summation Notation Expressions (continued)

```
62
63 *****
64 ** AI.1.3 SUM OF TWO ITEMS **
65 *****
66
67 PARAMETERS
68   SUM3A      SUM OF TWO ITEMS-1
69   SUM3B      SUM OF TWO ITEMS-1
70   A(I)       SUM OF TWO ITEMS-2
71   SUM4A      SUM OF TWO ITEMS-3
72   SUM4B      SUM OF TWO ITEMS-3;
73   SUM3A      = SUM(I, X(I)+SUM(J, Y(I, J)));
74   SUM3B      = SUM(I, X(I)) + SUM ((I,J), Y(I,J));
75   A(I)       = X(I) + SUM(J, Y(I,J));
76   SUM4A      = SUM(I, X(I)+Z(I));
77   SUM4B      = SUM(I, X(I)) + SUM(I, Z(I));
78   DISPLAY SUM3A, SUM3B, A, SUM4A, SUM4B;
79
80 *****
81 ** AI.2.1 FOR A SCALER EQUATION **
82 *****
83
84 PARAMETERS
85   B1      SUM FOR A SCALER EQUATION-1;
86   B1 = SUM((I,J,K), P(I,J,K)) + SUM((M,N), Q(M,N));
87   DISPLAY B1;
88
89 * $ONTEXT
90 * THE FOLLOWING SUMMATION NOTATIONS ARE INCORRECT
91 * IF YOU TURN THESE COMMANDS ON, YOU WILL ENCOUNTER
92 * ERROR MESSAGES
93 * PARAMETERS
94 *   B2      SUM FOR A SCALER EQUATION-2
95 *   B3      SUM FOR A SCALER EQUATION-3;
96 *   B2 = P(I,J,K) + Q(M,N);
97 *   B3 = SUM((I,J), P(I,J,K)) + SUM((M,N), Q(M,N));
98 *   DISPLAY B2, B3;
99 * $OFFTEXT
100
101 *****
102 ** A.I.2.2 FOR A FAMILY OF EQUATIONS **
103 *****
104
105 VARIABLES      C(I), D(I,J), E1(I,J), F(J);
106 EQUATIONS
107   EQC(I)      EQUATION C
108   EQD(I,J)    EQUATION D
109   EQE1(I,J)   EQUATION E1;
110   EQC(I)..    C(I) =E= 2;
111   EQD(I,J)..  D(I,J) =E= 2;
112   EQE1(I,J).. E1(I,J) =E= F(J);
113
114 * $ONTEXT
115 * THE FOLLOWING EXPRESSION IS INCORRECT
116 * ERROR MESSAGES WILL BE ENCOUNTERED
```

117 * VARIABLES E2(I,J);

Table 1. Sample GAMS Commands for Summation Notation Expressions (continued)

```
118 * EQUATION
119 *       EQE2      EQUATION E2;
120 *       EQE2..   E2(I,J) =E= 2;
121 * $OFFTEXT
122
123 VARIABLES G1(I), H1(I,J);
124 EQUATIONS
125       EQG1(I)    EQUATION G1
126       EQH1(I,J) EQUATION H1;
127       EQG1(I).. G1(I) =E= SUM((J,K), P(I,J,K));
128       EQH1(I,J).. H1(I,J) =E= SUM(K, P(I,J,K));
129
130 * $ONTEXT
131 * THE FOLLOWING EXPRESSIONS ARE INCORRECT
132 * ERROR MESSAGES WILL BE ENCOUNTERED
133 * VARIABLES G2(I), H2(I);
134 * EQUATIONS
135 *       EQG2(I)    EQUATION G2
136 *       EQH2(I)    EQUATION H2;
137 *       EQG2(I).. G2(I) =E= P(I,J,K);
138 *       EQH2(I).. H2(I) =E= SUM(K, P(I,J,K));
139 * $OFFTEXT
140
141 VARIABLES L1(I), U(I,M), R(I,J,K,M), S(I,J,M);
142 EQUATIONS
143       EQL1(I)    EQUATION L1
144       EQN(I,M)   EQUATION N;
145       EQL1(I).. L1(I) =E= SUM((J,K), P(I,J,K));
146       EQN(I,M).. U(I,M) =E= SUM((J,K),R(I,J,K,M)) + SUM(J, S(I,J,M));
147
148 * $ONTEXT
149 * THE FOLLOWING EXPRESSIONS ARE INCORRECT
150 * ERROR MESSAGES WILL BE ENCOUNTERED
151 * VARIABLES L2;
152 * EQUATIONS
153 *       EQL2(I)    EQUATION L2;
154 *       EQL2(I).. L2 =E= P(I,J,K);
155 * OFFTEXT
156
157 VARIABLE O(I,M);
158 EQUATION
159       EQO(I,M) EQUATION O;
160       EQO(I,M).. 2 + SUM((J,K), P(I,J,K)) + SUM(J, S(I,J,M)) =E=
O(I,M);
161
162
163 * $ONTEXT
164 * THE FOLLOWING EXPRESSION IS INCORRECT
165 * GAMS ERROR MESSAGES WILL BE ENCOUNTERED
166 * VARIABLE W(I);
167 * EQUATION
168 *       EQW(I)    EQUATION W;
169 *       EQW(I).. W(I) =E= SUM((I,J,K), P(I,J,K));
170 * $OFFTEXT
171
```

Table 1. Sample GAMS Commands for Summation Notation Expressions (continued)

```
172 *****
173 ** AI.4 DEFINING AND USING VARIABLES **
174 *****
175
176 VARIABLES
177     OBJ1          OBJECTIVE FUNCTION VALUE
178     T(I,J,K)     DECISION VARIABLE;
179 EQUATIONS
180     OBJFUNC1     OBJECTIVE FUNCTION
181     CONST(K)     CONSTRAINT;
182     OBJFUNC1..   OBJ1 =E= SUM((I,J,K), T(I,J,K));
183     CONST(K)..   SUM((I,J), T(I,J,K)) =E= 3;
184 MODEL EXAMPLE1 /ALL/;
185 SOLVE EXAMPLE1 USING LP MAXIMIZING OBJ1;
186 DISPLAY T.L;
187
188 * $ONTEXT
189 * THE FOLLOWING COMMANDS ARE INCORRECT
190 * THEY WILL RESULT IN ERROR MESSAGES
191 * VARIABLES
192 *     OBJ2          OBJECTIVE FUNCTION VALUE
193 *     TT(I,J,K)    DECISION VARIABLE;
194 * POSITIVE VARIABLE TT;
195 * EQUATIONS
196 *     OBJFUNC2     OBJECTIVE FUNCTION
197 *     CONSTT(K)    CONSTRAINT;
198 *     OBJFUNC2..   OBJ2 =E= SUM((I,J), TT(I,J));
199 *     CONSTT(K)..  SUM((I,J), TT(I,J,K)) =E= 3;
200 * MODEL EXAMPLE2 /ALL/;
201 * SOLVE EXAMPLE2 USING LP MAXIMIZING OBJ2;
202 * DISPLAY TT.L;
203 * $OFFTEXT
204
205 *****
206 ** AI.6 CAUTIONS AND EXTENSIONS **
207 *****
208
209 PARAMETER
210     SUM5A CAUTIONS AND EXTENSIONS-1;
211     SUM5A = SUM(J, X("2"));
212     DISPLAY SUM5A;
213
214 PARAMETER
215     PRODUCT6 CAUTIONS AND EXTENSIONS-2;
216     PRODUCT6 = PROD(I, X(I));
217     DISPLAY PRODUCT6;
218
219 PARAMETERS
220     SUM7A CAUTIONS AND EXTENSIONS-3
221     SUM7B CAUTIONS AND EXTENSIONS-3
222     SUM8A CAUTIONS AND EXTENSIONS-4
223     SUM8B CAUTIONS AND EXTENSIONS-4
224     SUM8C CAUTIONS AND EXTENSIONS-4
225     SUM9A CAUTIONS AND EXTENSIONS-5
```

Table 1. Sample GAMS Commands for Summation Notation Expressions (continued)

```

227 SUM10A CAUTIONS AND EXTENSIONS-6
228 SUM10B CAUTIONS AND EXTENSIONS-6;
229 SUM7A = SUM(I, 5*X(I));
230 SUM7B = 5*SUM(I, X(I));
231 SUM8A = SUM(I, 5*10);
232 SUM8B = 5*SUM(I, 10);
233 SUM8C = 5*3*10;
234 SUM9A = SUM((I,J), V(I,J)+Y(I,J));
235 SUM9B = SUM((I,J), V(I,J)) + SUM((I,J), Y(I,J));
236 SUM10A = SUM((I,J), X(I)+Y(I,J));
237 SUM10B = 2*SUM(I, X(I)) + SUM((I,J), Y(I,J));
238 DISPLAY SUM7A, SUM7B, SUM8A, SUM8B, SUM8C,
239 SUM9A, SUM9B, SUM10A, SUM10B;

```
